



Joshi, M., & Tulloh, R. (2017). Kawasaki disease and coronary artery aneurysms: From childhood to adulthood. *Future Cardiology*, 13(5), 491-501. <https://doi.org/10.2217/fca-2017-0039>

Peer reviewed version

Link to published version (if available):
[10.2217/fca-2017-0039](https://doi.org/10.2217/fca-2017-0039)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via [insert publisher name] at [insert hyperlink] . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

**Kawasaki Disease – Coronary artery aneurysms, from
Childhood to Adulthood**

Journal:	<i>Future Cardiology</i>
Manuscript ID	FCA-2017-0039
Manuscript Type:	Review
Keywords:	Kawasaki Disease, Coronary artery, Aneurysm

SCHOLARONE™
Manuscripts

Kawasaki Disease- coronary artery aneurysms

1
2
3
4
5
6
7
8
9
10
11
12
13
14

Abstract

Kawasaki disease (KD) is an acute, systemic vasculitis of childhood and confers a 25% risk of developing coronary artery aneurysms (CAA). Its aetiology is unknown, but the incidence is increasing rapidly with linked gene polymorphisms having been identified. A constellation of symptoms, epidemics and seasonality all implicate an unidentified infective or environmental cause. Intravenous immunoglobulin therapy (IVIG), aspirin and steroids all form the mainstay of acute treatment and reduces the incidence of CAA if given before 7 days. However, in some, these lesions persist and require ongoing management during follow up during childhood and into adult life. Evidence for further investigations in order to minimise complications is presented in order to minimise the myofibroblast proliferation and stenosis in the long term.

Keywords

Kawasaki disease, vasculitis, coronary artery, aneurysms.

Kawasaki Disease- coronary artery aneurysms

15

16 Background

17 Kawasaki disease (KD) is an acute, systemic vasculitis of medium-sized, primarily muscular arteries that typically
18 affects children and is generally self-limiting in nature(1). The incidence of KD highest in children under five
19 years, 85% of all diagnoses are made in this demographic, but can affect any age (2). It affects boys more than
20 girls, at a ratio of approximately 1.6:1 (3). The major complication is the development of coronary artery
21 aneurysms CAA; a potentially life-threatening condition. Consequently, there is a higher risk of myocardial
22 infarction, aneurysm rupture, coronary artery thrombus formation and sudden cardiac death.

23

24 The diagnostic criteria of complete KD are the presence of a fever lasting five days or more and at least four of
25 the following five criteria:

26

- 27 • bilateral, painless, non-exudative conjunctivitis
- 28 • cervical lymphadenopathy; generally larger than 1.5 cm and commonly unilateral
- 29 • polymorphous exanthema
- 30 • changes in lip or oral mucosa: commonly present as red, cracked lips; 'strawberry tongue' which is
31 seen as glossitis with hyperplastic fungiform papillae or generalised diffuse erythema or the oropharynx
- 32 • changes of the extremities, particularly erythema and oedema of the palms and soles of the feet; this
33 can progress later to desquamation of the skin

34

35 Incomplete KD may be seen with only 3 of the criteria in addition to fever; detection of coronary artery dilatation
36 or aneurysm on echocardiography requires only a fever and two of the criteria are to make a diagnosis of
37 atypical Kawasaki disease.

38

39 In developed countries, Kawasaki disease has overtaken rheumatic fever as the leading cause of acquired heart
40 disease in children (4). Rates of Kawasaki disease have steadily increased (5). In addition, while there is a
41 prevalence of Kawasaki disease in every country, Japan has the highest incidence of this condition and it is still
42 rising (6, 7).

Kawasaki Disease- coronary artery aneurysms

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69

The signs and symptoms of this condition develop over approximately 10 days and, in most cases, seemingly resolve spontaneously. However, this condition causes coronary artery aneurysms in up to 25% of children if left untreated: this is a statistic taken from the era of pre-intravenous immunoglobulin therapy, which is the mainstay treatment of Kawasaki disease today (8). Current data has shown a regression of coronary artery aneurysms in up to 75% of cases treated with IVIG(9), with approximately 3–6% of patients developing coronary artery aneurysms despite treatment (10, 11). 1% of patients develop giant coronary artery aneurysms, which carry the worst clinical prognosis.

There is also a subset of KD patients who fail to respond to IVIG; arguably demonstrating a more aggressive disease progression. The Kobayashi score uses laboratory findings and is routinely used in Japan to predict those with IVIG resistance, thus ensuring that second-line treatment is delivered promptly. However, this score did not predict response reliably in the US or the UK. Genetic variations between populations may lead to different laboratory results and therefore calculations may be unable to accurately predict IVIG resistance.

As a mechanism for the disease process remains unknown, no laboratory finding is pathognomonic for Kawasaki disease. Identification of KD is therefore dependent on clinical acumen alone. The constellation of symptoms is non-specific and is seen in other viral illnesses, which could delay prompt recognition and administration of treatment. Effective steps towards prevention are not feasible either and treatment options for the cardiac complications include invasive procedures such as angioplasties, stent placement and coronary-bypass surgeries (12). These reasons outline why Kawasaki disease is a paediatric research priority.

Aetiology

The aetiology of Kawasaki disease is poorly understood and several different theories have been proposed. Current evidence suggests the interaction of an unknown infective cause(s) (13) and a genetic predisposition are

Kawasaki Disease- coronary artery aneurysms

70 needed to develop KD. To what degree these factors influence KD, and if one is stronger than the other, has not
71 been formally elucidated.

72

73 It is, however, possible to argue that genetic factors have an important role in KD(14). Such genome wide
74 association studies have suggested that the polymorphism of the IgG receptor may have a part to play in the
75 susceptibility of children to KD and to the coronary artery aneurysms(15). The rates of Kawasaki disease are
76 significantly higher in East Asian populations: particularly Japan, which has the highest incidence of Kawasaki
77 disease in the world, at 264 per 100,000 children under 5 years (16). This is followed by Korea and Taiwan,
78 where rates are 134 and 66 respectively; the higher rates in these East Asian countries suggests some genetic
79 or environmental link. It could be argued that Kawasaki disease-associated pathogens (as yet undetermined)
80 show a degree of endemicity, or that the lifestyle followed in these three countries exposes children to the
81 pathogens, rather than any genetic bias conferring increased susceptibility. However, population studies in the
82 US show otherwise. The Japanese-American population that resides in Hawaii, USA has an incidence of 210 per
83 100,000, whereas the rate in Caucasian American children living in the same state is 13/100,000 (17). The
84 incidence in the Japanese-American population is approximately sixteen times greater than the Caucasian
85 population in the same environment, which could imply that the development of Kawasaki disease is dependent
86 on genetic factors and pathogen exposure (18).

87

88 Epidemiologically, the pattern of incidence suggests an infective cause. There have been three major nationwide
89 epidemics of Kawasaki disease in Japan, all demonstrating origination in a specific area and countrywide spread
90 over a 3-month period; similar to an epidemic (19). The acute, self-limiting nature of KD is another hallmark of
91 viral disease. Since the identification of Kawasaki disease, a multitude of different pathogens have been
92 investigated; these range from the bacterial (*Staphylococcus aureus*, *Klebsiella pneumoniae*, *Pseudomonas*
93 *aeruginosa*), to viral (herpesvirus, adenovirus and the Epstein-Barr virus) (20). 10% of patients with complete
94 Kawasaki disease have been shown to have low-level viral titres of human adenovirus present. The authors
95 concluded that this presented some link between infection with human adenovirus and the onset of Kawasaki
96 disease(21) but this has not yet been upheld as a definitive cause.

97

Kawasaki Disease- coronary artery aneurysms

However, it is important to note that titres of human adenovirus can persist years after the onset of the initial infection (22), and that this virus is particularly common in children: i.e. the 10% with detectable titres may have had the infection before, leading to the detectable viral load count. In addition, this study remarked that the sensitivity of their PCR quantification was high, but did not offer any way in which they could draw a discrepancy between viral load due to current infection or previous, and therefore unrelated, infection. Indeed, many papers have hypothesised different infective triggers; herpesvirus (20) is another example. The constellation of symptoms commonly seen in Kawasaki disease (fever, rash, irritability) are also seen in infection with adenoviruses and herpes viruses. Similar haematological abnormalities have also been observed, such as neutrophilia and high erythrocyte sedimentation rates (23). However, unlike the other two viruses mentioned, there is no evidence of human-to-human transmission in Kawasaki disease. Although this may appear to contradict with the earlier assertion of nationwide epidemics in Japan as these three events were comparable to viral outbreaks in their spread, epicentre, and seasonal timing, there was no evidence of human-to-human transmission found.

Peaks in occurrence appear to be dictated by the seasons, similar to what is seen in the pattern of viral infection (24). However, these peaks have a different seasonality across the world: while Japan, mainland US, Canada and the UK have all shown peaks during the winter months (25, 26) (27), in Hawaii however there was no seasonality at all, whereas China had its lowest incidence during the winter, and highest in the spring.

Peak incidence is in children between 18 and 24 months of age, with diagnoses of Kawasaki disease being made in children less than 3 months old being uncommon. This suggests that a child may have some protection from developing this disease until a certain age, implicating a role for transplacental antibodies. These antibodies play a powerful role in fighting bacterial infections, as demonstrated by children with congenital agammaglobulinemia (28) who commonly only begin to present with bacterial infections at 6 to 9 months (29); the onset of these bacterial infections do not generally coincide with the cessation of breast-feeding, implying that it is the antibodies transferred across the placental barrier that are responsible for this protective effect and not those found in the mother's breast milk. In addition, the effects of immunity conferred in utero tend to diminish

Kawasaki Disease- coronary artery aneurysms

within the first 3 months of life; further suggesting the role of transplacental antibodies. This pattern further suggests an infective cause being a trigger in developing Kawasaki disease.

By contrast, the rarity of Kawasaki disease after 5 years of age (30) implies that a matured immune system may be more adept at mounting an appropriate immune response: again, suggesting an infective cause. In addition, the greater incidence in boys supports this hypothesis as many infectious disease present more commonly in males than females (31). Furthermore, recurrence rates are low (at 3-4%), once again implying that the body is able to mount an immune response which prevents future cases (3).

Genetics

There is a greater risk of developing Kawasaki disease if there is an affected sibling or parent, implying a genetic role in the development of this disease (32). However, it is also important to note potential limitations of such data. (32). In this study, the number of patients affected with Kawasaki disease increased. The number of parents that had been diagnosed with KD in their childhood also increased. This trend displayed evidence of a genetic link between developing this condition. However, it is important to note that this study was carried out over a period of ten years from 1999 to 2008 and the increased proportion of affected patients reported every year may be due to an increased awareness of the condition amongst medical professionals, and therefore a greater rate of diagnosis, rather than an actual greater rate of prevalence.

In an attempt to understand the pathophysiology of Kawasaki disease, large genome studies have been conducted in populations where prevalence is high. Certain common single nucleotide polymorphisms have been observed in specific genes in individuals affected by KD. It is possible that these confer some level of genetic susceptibility in developing this condition. A functional polymorphism was identified that exceeded the formal threshold for genome-wide significance in the IgG receptor gene (*FCGR2A*) (15). Functional polymorphisms of this IgG receptor gene influence the phenotype of the individual by stratifying their responses to IgG as 'strong' or 'weak.' The particular single nucleotide polymorphism (SNP) that was identified had been shown to confer an increased risk of developing ulcerative colitis (33) in a Japanese population. This same SNP was shown to

Kawasaki Disease- coronary artery aneurysms

confer a decreased risk in developing systemic lupus erythematosus (34) in a European population. The involvement of this SNP in two autoimmune conditions suggests a dysfunctional immune response being fundamental in the development of Kawasaki disease in individuals. The role of an IgG-mediated immune response in the pathogenesis is also supported by IVIG being a mainstay of treatment, which will be discussed later in further detail.

Genomic studies showed three SNPs in a sodium-calcium transporter gene being validated for a significantly associated pathway (35). In KD patients, the carriers of homozygous 'risk' alleles of the *SLC8A1* gene have been shown to be at a greater risk of developing coronary artery aneurysms; the protein (NCX1) encoded was expressed in the inflammatory cells of the aneurysm wall. Altered calcium signalling appears to increase the risk of cardiac complications, implying a more aggressive course of disease. In addition, polymorphisms of *IPTKC*, a negative regulator of T-cell activations, have been associated with KD patients, and especially in those with coronary artery lesions (36). This is a Ca^{2+} /NFAT signalling pathway. Impaired calcium signalling suggests a more severe disease progression, as the functional polymorphism of *IPTKC* was strongly associated with patients who developed cardiac sequelae.

Calcineurin inhibitors have been shown to treat KD in IVIG-resistant patients (37): this suggests that impaired calcium signalling confers an increased risk of developing complications and its role should be further researched.

Up to 20% of KD patients fail to defervesce within 36 hours of IVIG therapy. It implies a more aggressive course of disease and longer period spent in acute inflammation, resulting in a greater likelihood of developing coronary artery aneurysm. The identification of these patients earlier could therefore improve clinical outcome, and so the Kobayashi score (38) was developed to predict patient response. It uses 7 laboratory and clinical findings to predict the unresponsiveness in KD patients to IVIG therapy. It has been used with positive results in the Japanese population; however, when used in the US and UK (39, 40), it failed to reliably predict IVIG responsiveness. Despite removal of patients with incomplete Kawasaki disease in the model, the Kobayashi

Kawasaki Disease- coronary artery aneurysms

score still failed to reliably predict IVIG response in a UK population. This suggests that the pattern of disease may differ according to ethnicity.

Pathophysiology

Although a specific infectious trigger has not yet been noted, the pathophysiology of this condition has been closely studied. Vasculitis has been observed in the cardiovascular, gastrointestinal, respiratory, integumentary, dermatological, urinary and nervous system(41); however, the most clinically significant is the vasculitis of coronary arteries, leading to coronary artery aneurysms. This is because it has resulted in myocardial infarction, ischaemic heart disease and sudden cardiac death in patients (42).

The progression of acute Kawasaki disease can be split into separate stages according to the number of days from the initial onset. Stage I (0-9 days after onset) is characterised by inflammation of the adventitia (also known as perivasculitis); stage II (12-25 days) often shows vasculitis of medium-sized, muscular arteries; stage III (28-31 days) is where pan-vasculitis is most prominent, and where arterial aneurysm development is commonly observed (3).

Analysis of tissue, blood and arterial walls has provided insight into the mechanism of disease progression. Tissue infiltration of lymphocytes, macrophages, plasma cells and large mononuclear cells demonstrates that there is an acute inflammatory response (43) and despite elevated levels of neutrophils in the blood (44) the lack of neutrophil invasion shows that these cells are not the chief mediator. While physiological levels of CD4 lymphocytes outnumber CD8, there was a 4–5 fold increase reported in CD8 lymphocytes in the arterial wall as compared to CD4. This is consistent with findings that are seen in the presence of an intracellular pathogen (45), thus providing evidence for a ubiquitous agent involved in the pathogenesis of Kawasaki disease. There was no significant change reported in the number of circulating CD8 lymphocytes in the acute stage, which indicates that the disease process actively recruits these lymphocytes into the arteries.

Kawasaki Disease- coronary artery aneurysms

An elevated IgA plasma cell proliferation (46) – greater than IgG and IgM – is seen in the arterial wall: this is unusual as IgA domination is seen in mucosal or lymphoid structures rather than vascular structures. This appears to suggest a stimulation of the immune system, likely at a mucosal site, which led to a systemic conversion of B lymphocytes to IgA secreting plasma cells.

A marked rise of IgA plasma cells appeared in the upper respiratory tract, pancreas, coronary arteries and kidneys of patients who had died of acute Kawasaki disease (47). This same study showed increased IgA plasma cell infiltration was also seen in children who had died of an acute respiratory tract infection; the similarity of these findings suggests the involvement of an infectious agent in Kawasaki disease. Studies have shown that the method by which IgA expansion is controlled is oligoclonal (46, 48), which suggests an infective cause. All synthetic antibodies that bound to acute KD tissues (48) detected antigen in all cytoplasmic inclusion bodies in acute KD affected ciliated bronchial epithelium. Inclusion bodies are pivotal in the pathophysiology and this finding is consistent with KD being caused by a microbial pathogen.

However, it is important to note that all of these studies used samples collected from deceased patients whose death was attributed to Kawasaki disease complications. Whilst all these data show similar results, which increases confidence in the conclusions obtained, it is unclear to what extent these conclusions can be extrapolated to patients that are a) alive, but with coronary artery lesions, b) alive, and with no coronary artery lesions and c) alive, with regressed coronary artery lesions. Due to the relatively recent history of Kawasaki disease, there has been little research into the coronary artery status of those who have died of a non-Kawasaki disease related cause, and therefore this still remains unknown. It will be interesting to study those particular patients; however, cataloguing and recruiting these patients will pose logistical challenges.

Treatment

Current guidelines outlining treatment of acute KD include intravenous immunoglobulin therapy and high-dose aspirin. Although aetiology and a definitive summary of the pathophysiology remain to be elucidated, numerous prospective, controlled trials have shown the efficacy of this regime (11, 49, 50).

Kawasaki Disease- coronary artery aneurysms

235

236 Intravenous immunoglobulin therapy in the treatment of KD has reduced rates of coronary artery aneurysm
237 formation from 25% to 3-6% in affected patients; however, around 20% of patients show IVIG-resistance, which
238 is qualified as a fever persisting 36 hours after IVIG was administered or a recrudescence fever (51, 52). C-
239 reactive protein levels persisting at >3 mg/L after 36 hours can also be a marker of IVIG-resistant disease.

240

241 2g/kg as a single dose is constituted as optimum therapy, and has shown greater efficacy than divided low-doses
242 (400 mg/kg/day) administered over several days. Meta-analysis concluded that both caused defervescence, but
243 the higher, single dose provided better protection against the formation of coronary artery lesions (49, 50). The
244 improvement of symptoms with IVIG and the association between a functional polymorphism in an IgG receptor
245 gene demonstrate that IgG has some role in the pathophysiology. In addition, even in IVIG-resistant patients, the
246 rash, swelling and erythema also associated with KD often improve (53). Reduced cytokine levels, decreased
247 number of circulating T cells, macrophages and neutrophils, and increased number of natural killer cells have all
248 been noted after IVIG treatment; however, how these changes are induced have not been described (53).

249

250 The use of aspirin remains slightly more controversial: the use of salicylates is strongly contraindicated in most
251 paediatric cases due to the risk of Reye's syndrome (54, 55) and KD is one of the few conditions in childhood
252 which is treated with aspirin. Although the duration of the characteristic fever appeared to be reduced by high-
253 dose aspirin therapy (56), there was no change noted in response to IVIG, resolution of inflammation, or
254 incidence of coronary artery lesion (57, 58). After the febrile period, aspirin therapy is reduced from 30-50
255 mg/kg/day to a low dose of 5mg/kg/day for 6 weeks and reassessed, depending on the presence or absence of
256 any coronary artery abnormalities.

257

258 IVIG-resistant therapy encompasses a second dose of IVIG at 2 g/kg, corticosteroids or both administered. Early
259 analyses showed a positive association between corticosteroid therapy and increased coronary artery
260 abnormalities. However, this finding has now been contributed to confounding, as in these retrospective studies
261 the sicker patients were more likely to be given corticosteroids, and therefore more likely to develop lesions (59).

262 The main indications for using adjunct corticosteroid therapy are:

Kawasaki Disease- coronary artery aneurysms

263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290

1. Patients who are IVIG resistant, with a persisting or recrudescant fever after 36 hours from IVIG treatment onset
2. Patients with features of very severe disease. In Japan, the Kobayashi score is used; however it has proven to be less accurate in other countries. Therefore, persistently high markers of cell injury and inflammation, such as raised CRP levels, liver dysfunction and anaemia are used
3. Patients with pre-existing coronary artery and/or peripheral aneurysms

Serum tumour necrosis factor- α (TNF- α) is elevated in KD patients, and in particular those who develop coronary artery aneurysms. Thus, anti TNF- α therapies such as infliximab are being investigated as alternative therapies; especially in patients unresponsive to IVIG. A 3-year-old patient, unresponsive to IVIG and methylprednisolone, defervesced in response to one dose of anti TNF- α (60). A retrospective study compared defervescence and length of hospital stay in IVIG-resistant patients treated with a second dose of IVIG and those treated with one dose of infliximab (61). Infliximab resulted in faster fever resolution and a shorter hospital stay. A recent randomised controlled trial (62) showed the same two results, with a greater response to infliximab (90.9%) than second-dose IVIG (65.6%). The size and power of these studies is limited, but demonstrates anti TNF- α as a potential therapeutic option.

Calcineurin inhibitors (such as cyclosporin A) have been used in the treatment of refractory KD (63, 64); however, they should be used with caution as they can be toxic to the endothelium (65). The decision to prescribe these medications must therefore be made on a case-by-case basis.

Coronary Artery Aneurysms

Current studies suggest that 20- 24% of children, even if treated, will suffer from coronary artery aneurysms(27). The definition of an aneurysm can now be recorded as an indexed (z score) size of the coronary artery, based on the body surface area of the child (66). Coronary artery dilation is defined as z score +2 to 2.5, small aneurysms are with z score \geq 2.5 but less than 5, moderate size CAA is seen in those with z score \geq 5 but less than 10 and those who have giant CAA have a z score \geq 10. We now know that it takes several years for many of these to

Kawasaki Disease- coronary artery aneurysms

291 resolve, with reducing likelihood of complete resolution correlating with increasing size and complexity of
292 coronary artery involvement (67). Recent AHA guidance details the long term outcome based on a severity
293 scoring system, which may be opinion for the USA, but which has little evidence base there and uncertain
294 relevance to the European or Asian population (68). We know that those children with giant CAA are those most
295 likely to suffer from Major Adverse Coronary Events in the longer term, necessitating antiplatelet and anti-
296 coagulant medication and multiple and regular repeat imaging.

297 For those children with just coronary dilation, or with no involvement at all, at the current time there is no
298 evidence from the western world that these children are at risk. There is evidence from a variety of studies that
299 the coronary arteries in this group may be abnormal in the long term in terms of coronary flow reserve (69) and
300 endothelial-dependent brachial artery reactivity (70) but there is dispute about the presence or absence of
301 coronary artery dysfunction (71) (72). No child, as far as we are aware, has been seen in the UK population,
302 without having had CAA, who has then grown up to demonstrate premature coronary artery disease. Hence the
303 recommendation is that these children can stop their aspirin and can be discharged from follow-up. The recent
304 AHA guidelines concur with this, but it is clear that until we have five decades of follow-up in the UK population,
305 we cannot be certain about the long-term outcome in this group. Since this group is 76% of the current UK
306 incident population of children with KD (27), it will be important to revisit this in future years when we have longer
307 term follow up.

308
309 There is some evidence that those children who had small or medium CAA, in whom these resolved, should be
310 followed up in the longer term (68, 73, 74). Our group have shown that they have circulating endothelial cells
311 (75). This is in line with the American Heart Association Guidelines (76) (68) that suggest that there may be
312 ongoing risk in these patients and advise ongoing imaging and dynamic myocardial and coronary flow studies.
313 Long term outcome after even moderate aneurysms suggests that there may be coronary artery stenosis, only
314 recognised by thallium scanning and not by stress testing (77). Hence there is recommendation that these
315 children can either stop or continue their aspirin and could have long-term follow-up. In the UK, there is no
316 evidence, yet, that these children are at risk of coronary events

Kawasaki Disease- coronary artery aneurysms

The evidence that those children with persistent CAA may develop complications is now overwhelming. The risk of Major Adverse Coronary Event (MACE) increases with size, complexity and location of CAA (9) (68). In those with persisting CAA >4mm, there is a high risk of coronary artery endothelial thickening which might lead to complications of ischaemia, especially on exercise (78). In those with giant CAA (>8mm) or z>10, long-term outcome is poor, with 40% having MACE with myocardial infarction often leading to death.

Kawasaki disease in Adulthood

There has been no formal collection of data regarding the number of adult patients with a previous history of Kawasaki disease in the US or the UK. Estimations are that approximately 24,000 young adults (≥ 18 years) in the US (42), with 8,400 of these patients classified at an AHA Risk level II or higher. Estimates for the UK have not yet been published but it appears that about 4% of young adults presenting with myocardial infarction have coronary angiograms indicative of previous KD (79). It is accepted that those patients who presented with giant aneurysms should undergo regular and routine follow up through adulthood. However, it is unclear what level of follow up treatment should be given to those with regressed aneurysms and those who presented with no coronary artery abnormalities. Prompt treatment with IVIG means that the majority of patients (76%) do not have these coronary artery lesions in the acute stage of KD.

Due to the growing rate of diagnosis, cardiologists will be dealing with adult patients who have long-term sequelae from this condition. Although coronary artery aneurysms are the main complication, other cardiovascular abnormalities should also be investigated: myocarditis and valvulitis (80) has also been observed in KD patients. Myocardial fibrosis and aortic and mitral regurgitation can follow and lead to death.

Studies in Japan showed that the standard mortality ratio (SMR) for young males with previous KD and cardiac sequelae was significantly higher than females with KD and age-matched population (81). This is one of few studies conducted that has investigated mortality rates in people with KD with no cardiac sequelae, and may indicate that patients with KD and no sequelae do not need routine follow-up. It suggests that paediatricians can

Kawasaki Disease- coronary artery aneurysms

stop seeing these patients during childhood, as this demographic does not have a significantly greater risk of mortality.

Childhood diagnosis of KD has been cited as a risk factor for accelerated coronary artery disease in adulthood (82) irrespective of lipid profile, suggesting that these patients should be followed up by cardiologists routinely. The only patients included in the test group for this study were patients with a historical KD diagnosis and cardiac sequelae: whether a KD diagnosis alone is a risk factor for accelerated atherosclerosis cannot be concluded. KD being a risk factor can be cited as a reason to follow up all patients in later life. However, as only those with coronary artery lesions were included in this study, it is not entirely possible to state that KD with no cardiac sequelae is also a risk factor.

Autopsy results of those with KD who died after the acute stage (all individuals were ≥ 15 years old at the time of death) showed different histological results in non-aneurysm coronary arteries(83). There was increased 'new' intimal thickening of the non-affected arteries, along with previous thickening during the arteritis phase in acute KD. Although these arteries were declared non-affected, there were still histological changes that remained during the patient's life, and can therefore not be excluded as a risk factor for atherosclerosis. The coronary arteries of a KD patient with no coronary artery lesion were compared to KD patients with the cardiac sequelae, and both showed platelet-derived growth factor (A) and inducible nitrate synthase in intimal smooth muscle cells (84). This was not seen in patients with no KD. Inflammatory markers were still present in this patient at 13 months post diagnosis, with an apparently normal coronary artery, and therefore KD with no coronary artery lesions can still pose as a risk factor for atherosclerosis. Although these are the histological findings of only one patient, and the results cannot be extrapolated to all, autopsy reports of children with no cardiac sequelae are incredibly rare. Obtaining tissue from patients to investigate this poses a challenge in itself, and in light of the data available, it is better practice to regularly follow up KD patients: even those with no history of coronary artery aneurysms.

Future perspective

Kawasaki Disease- coronary artery aneurysms

There are a variety of issues yet to be solved. First of all we need a certain marker for diagnosis and a known aetiology. Then we need to have better recognition and diagnosis with more effective acute treatment to reduce the incidence of coronary artery aneurysms. This would reduce long term complications and would be best accompanied by effective therapy at reducing myofibroblastic proliferation and hence the risk of major adverse coronary events.

Executive summary

In summary, therefore, we have described the presentation, treatment and long term management strategy for children and young adults with this increasingly common disease.

Financial Disclosure

Neither author has any financial disclosures relevant to this manuscript. No commercial involvement was connected to this study at any stage. Both authors contributed to the writing.

References

Uncategorized References

1. Eleftheriou D, Levin M, Shingadia D, Tulloh R, Klein NJ, Brogan PA. Management of Kawasaki disease. Archives of disease in childhood. 2014;99(1):74-83.
2. Pinna GS, Kafetzis DA, Tselkas OI, Skevaki CL. Kawasaki disease: an overview. Current opinion in infectious diseases. 2008;21(3):263-70.
3. Burns JC, Kushner HI, Bastian JF, Shike H, Shimizu C, Matsubara T, et al. Kawasaki disease: A brief history. Pediatrics. 2000;106(2):E27.
4. Daniels LB, Tjajadi MS, Walford HH, Jimenez-Fernandez S, Trofimenko V, Fick DB, et al. Prevalence of Kawasaki Disease in Young Adults with Suspected Myocardial Ischemia. Circulation. 2012;125(20):2447-53.
5. Singh S, Vignesh P, Burgner D. The epidemiology of Kawasaki disease: a global update. Archives of disease in childhood. 2015;100(11):1084-8.
6. Nakamura Y, Yashiro M, Uehara R, Sadakane A, Tsuboi S, Aoyama Y, et al. Epidemiologic features of Kawasaki disease in Japan: results of the 2009-2010 nationwide survey. J Epidemiol. 2012;22(3):216-21.
7. Nakamura Y, Yashiro M, Uehara R, Oki I, Kayaba K, Yanagawa H. Increasing incidence of Kawasaki disease in Japan: nationwide survey. Pediatrics international : official journal of the Japan Pediatric Society. 2008;50(3):287-90.
8. Kato H, Sugimura T, Akagi T, Sato N, Hashino K, Maeno Y, et al. Long-term consequences of Kawasaki disease. A 10- to 21-year follow-up study of 594 patients. Circulation. 1996;94(6):1379-85.

Kawasaki Disease- coronary artery aneurysms

9. Friedman KG, Gauvreau K, Hamaoka-Okamoto A, Tang A, Berry E, Tremoulet AH, et al. Coronary Artery Aneurysms in Kawasaki Disease: Risk Factors for Progressive Disease and Adverse Cardiac Events in the US Population. *Journal of the American Heart Association*. 2016;5(9).
10. Ogata S, Tremoulet AH, Sato Y, Ueda K, Shimizu C, Sun X, et al. Coronary artery outcomes among children with Kawasaki disease in the United States and Japan. *International journal of cardiology*. 2013;168(4):3825-8.
11. Newburger JW, Takahashi M, Burns JC, Beiser AS, Chung KJ, Duffy CE, et al. The treatment of Kawasaki syndrome with intravenous gamma globulin. *The New England journal of medicine*. 1986;315(6):341-7.
12. Akagi T, Ogawa S, Ino T, Iwasa M, Echigo S, Kishida K, et al. Catheter interventional treatment in Kawasaki disease: A report from the Japanese Pediatric Interventional Cardiology Investigation group. *The Journal of pediatrics*. 2000;137(2):181-6.
13. Burns JC, Glode MP. Kawasaki syndrome. *Lancet*. 2004;364(9433):533-44.
14. Khor CC, Davila S, Shimizu C, Sheng S, Matsubara T, Suzuki Y, et al. Genome-wide linkage and association mapping identify susceptibility alleles in ABCC4 for Kawasaki disease. *Journal of medical genetics*. 2011;48(7):467-72.
15. Khor CC, Davila S, Breunis WB, Lee YC, Shimizu C, Wright VJ, et al. Genome-wide association study identifies FCGR2A as a susceptibility locus for Kawasaki disease. *Nature genetics*. 2011;43(12):1241-6.
16. Harnden A, Tulloh R, Burgner D. Kawasaki disease. *Bmj*. 2014;349:g5336.
17. Uehara R, Belay ED. Epidemiology of Kawasaki Disease in Asia, Europe, and the United States. *J Epidemiol*. 2012. p. 79-85.
18. Rodo X, Curcoll R, Robinson M, Ballester J, Burns JC, Cayan DR, et al. Tropospheric winds from northeastern China carry the etiologic agent of Kawasaki disease from its source to Japan. *Proceedings of the National Academy of Sciences of the United States of America*. 2014;111(22):7952-7.
19. Yanagawa H, Nakamura Y, Ojima T, Yashiro M, Tanihara S, Oki I. Changes in epidemic patterns of Kawasaki disease in Japan. *The Pediatric infectious disease journal*. 1999;18(1):64-6.
20. Shingadia D, Bose A, Booy R. Could a herpesvirus be the cause of Kawasaki disease? *The Lancet infectious diseases*. 2002;2(5):310-3.
21. Jaggi P, Kajon AE, Mejias A, Ramilo O, Leber A. Human adenovirus infection in Kawasaki disease: a confounding bystander? *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2013;56(1):58-64.
22. Garnett CT, Erdman D, Xu W, Gooding LR. Prevalence and quantitation of species C adenovirus DNA in human mucosal lymphocytes. *J Virol*. 2002;76(21):10608-16.
23. Larranaga C, Kajon A, Villagra E, Avendano LF. Adenovirus surveillance on children hospitalized for acute lower respiratory infections in Chile (1988-1996). *J Med Virol*. 2000;60(3):342-6.
24. Fisman D. Seasonality of viral infections: mechanisms and unknowns. *Clin Microbiol Infect*. 2012;18(10):946-54.
25. Park YW, Han JW, Hong YM, Ma JS, Cha SH, Kwon TC, et al. Epidemiological features of Kawasaki disease in Korea, 2006-2008. *Pediatr Int*. 2011;53(1):36-9.
26. Huang WC, Huang LM, Chang IS, Chang LY, Chiang BL, Chen PJ, et al. Epidemiologic features of Kawasaki disease in Taiwan, 2003-2006. *Pediatrics*. 2009;123(3):e401-5.
27. Tulloh RM, Ramanan A, Brogan P, Tizard EJ, Connolly GM, Harnden A, et al. Kawasaki disease. Results of the BPSU survey in UK and Ireland. *Archives of disease in childhood*. 2016;101:suppl 1, A5.
28. Winkelstein JA, Marino MC, Lederman HM, Jones SM, Sullivan K, Burks AW, et al. X-linked agammaglobulinemia: report on a United States registry of 201 patients. *Medicine (Baltimore)*. 2006;85(4):193-202.

Kawasaki Disease- coronary artery aneurysms

29. Gedalia A. Kawasaki disease: 40 years after the original report. *Current rheumatology reports*. 2007;9(4):336-41.
30. Van Camp G, Deschamps P, Mestrez F, Levy J, Van Laethem Y, de Marneffe M, et al. Adult onset Kawasaki disease diagnosed by the echocardiographic demonstration of coronary aneurysms. *European heart journal*. 1995;16(8):1155-7.
31. Green MS. The male predominance in the incidence of infectious diseases in children: a postulated explanation for disparities in the literature. *Int J Epidemiol*. 1992;21(2):381-6.
32. Uehara R, Yashiro M, Nakamura Y, Yanagawa H. Parents with a history of Kawasaki disease whose child also had the same disease. *Pediatrics international : official journal of the Japan Pediatric Society*. 2011;53(4):511-4.
33. Asano K, Matsushita T, Umeno J, Hosono N, Takahashi A, Kawaguchi T, et al. A genome-wide association study identifies three new susceptibility loci for ulcerative colitis in the Japanese population. *Nat Genet*. 2009;41(12):1325-9.
34. Harley JB, Alarcon-Riquelme ME, Criswell LA, Jacob CO, Kimberly RP, Moser KL, et al. Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in ITGAM, PTK, KIAA1542 and other loci. *Nat Genet*. 2008;40(2):204-10.
35. Shimizu C, Eleftherohorinou H, Wright VJ, Kim J, Alphonse MP, Perry JC, et al. Genetic Variation in the SLC8A1 Calcium Signaling Pathway Is Associated With Susceptibility to Kawasaki Disease and Coronary Artery Abnormalities. *Circ Cardiovasc Genet*. 2016;9(6):559-68.
36. Onouchi Y, Gunji T, Burns JC, Shimizu C, Newburger JW, Yashiro M, et al. ITPKC functional polymorphism associated with Kawasaki disease susceptibility and formation of coronary artery aneurysms. *Nat Genet*. 2008;40(1):35-42.
37. Tremoulet AH. Calcineurin Inhibitor Treatment of Intravenous Immunoglobulin-Resistant Kawasaki Disease. 2012;161(3):506-12.e1.
38. Kobayashi T, Inoue Y, Takeuchi K, Okada Y, Tamura K, Tomomasa T, et al. Prediction of intravenous immunoglobulin unresponsiveness in patients with Kawasaki disease. *Circulation*. 2006;113(22):2606-12.
39. Sleeper LA, Minich LLA, McCrindle BM, Li JS, Mason W, Colan SD, et al. EVALUATION OF KAWASAKI DISEASE RISK SCORING SYSTEMS FOR INTRAVENOUS IMMUNOGLOBULIN RESISTANCE. *J Pediatr*. 2011;158(5):831-5.e3.
40. Davies S, Sutton N, Blackstock S, Gormley S, Hoggart CJ, Levin M, et al. Predicting IVIG resistance in UK Kawasaki disease. *Archives of disease in childhood*. 2015;100(4):366-8.
41. Takahashi K, Oharaseki T, Yokouchi Y. Update on etio and immunopathogenesis of Kawasaki disease. *Current opinion in rheumatology*. 2014;26(1):31-6.
42. Gordon JB, Kahn AM, Burns JC. When children with Kawasaki disease grow up: Myocardial and vascular complications in adulthood. *J Am Coll Cardiol*. 2009;54(21):1911-20.
43. Rowley AH, Baker SC, Orenstein JM, Shulman ST. Searching for the cause of Kawasaki disease--cytoplasmic inclusion bodies provide new insight. *Nature reviews Microbiology*. 2008;6(5):394-401.
44. Amano S, Hazama F, Kubagawa H, Tasaka K, Haebara H, Hamashima Y. General pathology of Kawasaki disease. On the morphological alterations corresponding to the clinical manifestations. *Acta pathologica japonica*. 1980;30(5):681-94.
45. Brown TJ, Crawford SE, Cornwall ML, Garcia F, Shulman ST, Rowley AH. CD8 T lymphocytes and macrophages infiltrate coronary artery aneurysms in acute Kawasaki disease. *The Journal of infectious diseases*. 2001;184(7):940-3.
46. Rowley AH, Shulman ST, Spike BT, Mask CA, Baker SC. Oligoclonal IgA response in the vascular wall in acute Kawasaki disease. *Journal of immunology*. 2001;166(2):1334-43.
47. Rowley AH, Shulman ST, Mask CA, Finn LS, Terai M, Baker SC, et al. IgA plasma cell infiltration of proximal respiratory tract, pancreas, kidney, and coronary artery in acute Kawasaki disease. *The Journal of infectious diseases*. 2000;182(4):1183-91.

Kawasaki Disease- coronary artery aneurysms

48. Rowley AH, Shulman ST, Garcia FL, Guzman-Cottrill JA, Miura M, Lee HL, et al. Cloning the arterial IgA antibody response during acute Kawasaki disease. *Journal of immunology*. 2005;175(12):8386-91.
49. Newburger JW, Takahashi M, Beiser AS, Burns JC, Bastian J, Chung KJ, et al. A single intravenous infusion of gamma globulin as compared with four infusions in the treatment of acute Kawasaki syndrome. *The New England journal of medicine*. 1991;324(23):1633-9.
50. Durongpisitkul K, Gururaj VJ, Park JM, Martin CF. The prevention of coronary artery aneurysm in Kawasaki disease: a meta-analysis on the efficacy of aspirin and immunoglobulin treatment. *Pediatrics*. 1995;96(6):1057-61.
51. Hashino K, Ishii M, Iemura M, Akagi T, Kato H. Re-treatment for immune globulin-resistant Kawasaki disease: a comparative study of additional immune globulin and steroid pulse therapy. *Pediatrics international : official journal of the Japan Pediatric Society*. 2001;43(3):211-7.
52. Wallace CA, French JW, Kahn SJ, Sherry DD. Initial intravenous gammaglobulin treatment failure in Kawasaki disease. *Pediatrics*. 2000;105(6):E78.
53. Burns JC, Franco A. The immunomodulatory effects of intravenous immunoglobulin therapy in Kawasaki disease. *Expert review of clinical immunology*. 2015;11(7):819-25.
54. Macdonald S. Aspirin use to be banned in under 16 year olds. *Bmj*. 2002;325(7371):988.
55. Belay ED, Bresee JS, Holman RC, Khan AS, Shahriari A, Schonberger LB. Reye's syndrome in the United States from 1981 through 1997. *N Engl J Med*. 1999;340(18):1377-82.
56. Lee G, Lee SE, Hong YM, Sohn S. Is high-dose aspirin necessary in the acute phase of kawasaki disease? *Korean circulation journal*. 2013;43(3):182-6.
57. Akagi T, Kato H, Inoue O, Sato N. A study on the optimal dose of aspirin therapy in Kawasaki disease--clinical evaluation and arachidonic acid metabolism. *The Kurume medical journal*. 1990;37(3):203-8.
58. Terai M, Shulman ST. Prevalence of coronary artery abnormalities in Kawasaki disease is highly dependent on gamma globulin dose but independent of salicylate dose. *The Journal of pediatrics*. 1997;131(6):888-93.
59. Kato H, Koike S, Yokoyama T. Kawasaki disease: effect of treatment on coronary artery involvement. *Pediatrics*. 1979;63(2):175-9.
60. Weiss JE, Eberhard BA, Chowdhury D, Gottlieb BS. Infliximab as a novel therapy for refractory Kawasaki disease. *The Journal of rheumatology*. 2004;31(4):808-10.
61. Son MB, Gauvreau K, Burns JC, Corinaldesi E, Tremoulet AH, Watson VE, et al. Infliximab for intravenous immunoglobulin resistance in Kawasaki disease: a retrospective study. *The Journal of pediatrics*. 2011;158(4):644-9 e1.
62. Youn Y, Kim J, Hong YM, Sohn S. Infliximab as the First Retreatment in Patients with Kawasaki Disease Resistant to Initial Intravenous Immunoglobulin. *The Pediatric infectious disease journal*. 2016;35(4):457-9.
63. Suzuki H, Terai M, Hamada H, Honda T, Suenaga T, Takeuchi T, et al. Cyclosporin A treatment for Kawasaki disease refractory to initial and additional intravenous immunoglobulin. *The Pediatric infectious disease journal*. 2011;30(10):871-6.
64. Hamada H, Suzuki H, Abe J, Suzuki Y, Suenaga T, Takeuchi T, et al. Inflammatory cytokine profiles during Cyclosporin treatment for immunoglobulin-resistant Kawasaki disease. *Cytokine*. 2012;60(3):681-5.
65. Akman-Demir G, Ayranci O, Kurtuncu M, Vanli EN, Mutlu M, Tugal-Tutkun I. Cyclosporine for Behcet's uveitis: is it associated with an increased risk of neurological involvement? *Clin Exp Rheumatol*. 2008;26(4 Suppl 50):S84-90.
66. McCrindle BW, Li JS, Minich LL, Colan SD, Atz AM, Takahashi M, et al. Coronary artery involvement in children with Kawasaki disease: risk factors from analysis of serial normalized measurements. *Circulation*. 2007;116(2):174-9.
67. Hillary C JD, Khader Y, Tulloh R. How long does it take for coronary artery aneurysms to resolve after Kawasaki disease? *Cardiology in the young*. 2017;27(Supplement S2):S42.

Kawasaki Disease- coronary artery aneurysms

68. McCrindle BW, Rowley AH, Newburger JW, Burns JC, Bolger AF, Gewitz M, et al. Diagnosis, Treatment, and Long-Term Management of Kawasaki Disease: A Scientific Statement for Health Professionals From the American Heart Association. *Circulation*. 2017.
69. Muzik O, Paridon SM, Singh TP, Morrow WR, Dayanikli F, Di Carli MF. Quantification of myocardial blood flow and flow reserve in children with a history of Kawasaki disease and normal coronary arteries using positron emission tomography. *Journal of the American College of Cardiology*. 1996;28(3):757-62.
70. Dhillon R, Clarkson P, Donald AE, Powe AJ, Nash M, Novelli V, et al. Endothelial dysfunction late after Kawasaki disease. *Circulation*. 1996;94(9):2103-6.
71. Mitani Y, Okuda Y, Shimpo H, Uchida F, Hamanaka K, Aoki K, et al. Impaired endothelial function in epicardial coronary arteries after Kawasaki disease. *Circulation*. 1997;96(2):454-61.
72. Yamakawa R, Ishii M, Sugimura T, Akagi T, Eto G, Iemura M, et al. Coronary endothelial dysfunction after Kawasaki disease: evaluation by intracoronary injection of acetylcholine. *Journal of the American College of Cardiology*. 1998;31(5):1074-80.
73. Iemura M, Ishii M, Sugimura T, Akagi T, Kato H. Long term consequences of regressed coronary aneurysms after Kawasaki disease: vascular wall morphology and function. *Heart*. 2000;83(3):307-11.
74. Dietz SM, Tacke CE, Kuipers IM, Wiegman A, de Winter RJ, Burns JC, et al. Cardiovascular imaging in children and adults following Kawasaki disease. *Insights into imaging*. 2015;6(6):697-705.
75. Shah V, Christov G, Mukasa T, Brogan KS, Wade A, Eleftheriou D, et al. Cardiovascular status after Kawasaki disease in the UK. *Heart*. 2015;101(20):1646-55.
76. Newburger JW, Takahashi M, Gerber MA, Gewitz MH, Tani LY, Burns JC, et al. Diagnosis, treatment, and long-term management of Kawasaki disease: a statement for health professionals from the Committee on Rheumatic Fever, Endocarditis and Kawasaki Disease, Council on Cardiovascular Disease in the Young, American Heart Association. *Circulation*. 2004;110(17):2747-71.
77. Fukushima J, Takahashi N, Ueda K, Hijii T, Igarashi H, Ohshima A. Long-term outcome of coronary abnormalities in patients after Kawasaki disease. *Pediatric cardiology*. 1996;17(2):71-6.
78. Tsuda E, Kamiya T, Kimura K, Ono Y, Echigo S. Coronary artery dilatation exceeding 4.0 mm during acute Kawasaki disease predicts a high probability of subsequent late intima-medial thickening. *Pediatric cardiology*. 2002;23(1):9-14.
79. Waterhouse B, Tulloh RM, Johnson T. Prevalence of coronary artery aneurysms in patients under the age of 50 presenting with STEMI : evaluating the impact of untreated Kawasaki disease. *European heart journal*. 2016;37:Suppl 1, 191.
80. Akagi T, Kato H, Inoue O, Sato N, Imamura K. Valvular heart disease in Kawasaki syndrome: incidence and natural history. *American heart journal*. 1990;120(2):366-72.
81. Nakamura Y, Aso E, Yashiro M, Uehara R, Watanabe M, Oki I, et al. Mortality among persons with a history of kawasaki disease in Japan: mortality among males with cardiac sequelae is significantly higher than that of the general population. *Circulation journal : official journal of the Japanese Circulation Society*. 2008;72(1):134-8.
82. Noto N, Okada T, Yamasuge M, Taniguchi K, Karasawa K, Ayusawa M, et al. Noninvasive assessment of the early progression of atherosclerosis in adolescents with Kawasaki disease and coronary artery lesions. *Pediatrics*. 2001;107(5):1095-9.
83. Takahashi K, Oharaseki T, Naoe S. Pathological study of postcoronary arteritis in adolescents and young adults: with reference to the relationship between sequelae of Kawasaki disease and atherosclerosis. *Pediatric cardiology*. 2001;22(2):138-42.
84. Suzuki A, Miyagawa-Tomita S, Komatsu K, Nakazawa M, Fukaya T, Baba K, et al. Immunohistochemical study of apparently intact coronary artery in a child after Kawasaki disease. *Pediatrics international : official journal of the Japan Pediatric Society*. 2004;46(5):590-6.